


The arithmetic of linear orders

Garrett Ervin
joint with Eric Paul

June 5, 2024

A puzzle



Is C isomorphic to D ?

Goal: Study the arithmetic of the class of linear orders LO under the sum $+$ and lexicographic product \times .

- ▶ Arithmetic of $(LO, +)$ worked out classically by Tarski, Aronszajn, and especially Lindenbaum.
- ▶ Much less known about arithmetic in (LO, \times) ; lone classical result due to Morel characterizing cancellation on the right.
- ▶ We give some new results concerning cancellation on the left.

Defining the sum and product

Definition: Given linear orders A and B :

- ▶ The *sum* $A + B$ is the order obtained by placing a copy of B to the right of A (“ A followed by B ”),
- ▶ The *lexicographic product* $A \times B = AB$ is the order obtained by replacing every point in A with a copy of B (“ A -many copies of B ”).

Example: If

$$\begin{array}{lcl} A & = & \bullet \bullet \bullet \\ B & = & \text{—————} \end{array}$$

Then

$$\begin{array}{lcl} A + B & = & \bullet \bullet \bullet \text{ —————} \\ A \times B & = & \text{—————} \text{ —————} \text{ —————} \end{array}$$

Some examples

$$\omega + 1 = \text{purple dots} \dots \text{blue dot} \not\cong \omega$$

$$1 + \omega = \text{blue dot} \text{purple dots} \dots \cong \omega$$

$$2\mathbb{Z} = (\text{purple dots}) (\text{purple dots}) \not\cong \mathbb{Z}$$

$$\mathbb{Z}^2 = \text{purple dots} \text{purple dots} \dots \cong \mathbb{Z}$$

$$2\mathbb{Q} = (\text{purple dots}) (\text{purple dots}) \cong \mathbb{Q}$$

$$\mathbb{Q}^2 = (\text{purple dots}) (\text{purple dots}) (\text{purple dots}) (\text{purple dots}) \not\cong \mathbb{Q}$$

Arithmetic of $(LO, +)$

Question: to what extent do familiar laws of $(\mathbb{N}, +)$ hold in $(LO, +)$?

- ▶ E.g. the additive cancellation law, unique division by n , commutativity.
- ▶ Results due to Tarski, Aronszajn, and especially Lindenbaum.




A. Lindenbaum (1904-1941)

Arithmetic of $(LO, +)$

To motivate the results, let's consider simple “equations” (i.e. isomorphisms) over LO involving $+$.


We begin with the three-term isomorphism $A + B \cong C$:



If we add constraints by setting certain terms equal, we get a recurrence that we can then attempt to “solve.”

Left absorption

Consider $A + X \cong X$:





Always true if $A = 0$. But can have $A \neq 0$:

- ▶ E.g. if $A = 1$ and $X = 1 + 1 + \dots = \omega$.
- ▶ More generally, if A is arbitrary and $X = A + A + \dots = \omega A$.
- ▶ More generally still, if A, R are arbitrary and $X = \omega A + R$.

Left absorption

Thm (folklore): If $A + X \cong X$, then $X \cong \omega A + R$ for some R .

Proof:



$$X \cong \omega A + R$$

Absorption implies non-cancellation

If $A + X \cong X$ and $A \not\cong 0$, then X cannot be cancelled in the isomorphism $A + X \cong X$.

- ▶ So, right cancellation fails in $(LO, +)$.
- ▶ But, for this form of non-right-cancellation (left absorption), we can completely characterize the failure.


Symmetrically, we can show $X + A \cong X$ iff $X \cong L + \omega^*A$.

- ▶ Left cancellation fails too...

Can we characterize the solutions of $X + X \cong X$? Stay tuned!

More arithmetic in $(LO, +)$


Now let's consider the four-term isomorphism $A + B \cong C + D$:



We get a number of familiar recurrences from this isomorphism by setting terms equal.

Cancelling on the right

Consider $A + X \cong B + X$:



Can we cancel X and conclude $A \cong B$? Not in general.


- ▶ E.g. if $A \neq 0$, $B = 0$, and X absorbs A on the left.
- ▶ *It turns out:* left absorption is the *only* barrier to right cancellation.

X right cancels $\Leftrightarrow X$ does not left absorb

Thm (folklore): If $A + X \cong B + X$, then either $A \cong B$ or there is a non-empty order K such $K + X \cong X$.

($X + A \cong X + B$ is symmetric.)


Proof:



Another view: if $A + X \cong B + X$, then A and B are *almost* isomorphic (up to a “negligible” final segment absorbed by X).

Dividing by 2

Now suppose $X + X \cong Y + Y$:



Does it follow $X \cong Y$?


Dividing by 2

Thm (Lindenbaum): If X is isomorphic to a final segment of Y and Y is isomorphic to an initial segment of X , then $X \cong Y$.

Proof: Cantor-Schroeder-Bernstein proof works!

Cor (Lindenbaum): If $X + X \cong Y + Y$ then $X \cong Y$.


Proof:



Dividing by n

More generally we have:

Thm (Lindenbaum): if $nX \cong nY$ then $X \cong Y$.



► Proof harder for $n > 2$.

Fractions

What if $nX \cong mY$ with $n \neq m$?


- ▶ By cancelling common factors, suffices to assume $\gcd(n, m) = 1$.

Thm (Lindenbaum): If $nX \cong mY$ with $\gcd(n, m) = 1$, then there is a linear order C such that $X \cong mC$ and $Y \cong nC$.

E.g. if $2X \cong 3Y$, then $\exists C$ s.t. $X \cong 3C$ and $Y \cong 2C$.

Fractions

Recall our puzzle: is C isomorphic to D ?



...the answer is yes!

The proof of Lindenbaum's theorem is tricky. It cases out over a fundamental dichotomy:


Thm (Lindenbaum, Tarski): For a linear order X , exactly one holds:

- i. $mX \not\cong nX$ for all $m, n \in \mathbb{N}$ with $m \neq n$,
- ii. $mX \cong nX$ for all $m, n \geq 1$.

i.e., the finite multiples of a linear order X are either all distinct or all the same.

Commuting pairs

Now let's consider solutions to the isomorphism $X + Y \cong Y + X$:



There are two “obvious” ways the isomorphism can hold:

- i. (finite sum) $\exists C$ s.t. $X \cong nC$ and $Y \cong mC$ for some $m, n \in \mathbb{N}$,
- ii. (bi-absorption) $X + Y \cong Y + X \cong Y$.


Commuting pairs

Thm (Tarski): These are the only ways if X, Y are countable or if X, Y are scattered.

Conj (Tarski): These are the only ways for any linear orders X, Y .

Prop'n (Lindenbaum): There is another way.

Thm (Aronszajn): There is only one other way.



Arithmetic of $(LO, +)$: summary

$$A + X \cong X \quad \text{iff} \quad X \cong \omega A + R \quad (A \text{ "almost } \cong" 0)$$

($X + A \cong X$ symmetric)

$$A + X \cong B + X \quad \text{iff} \quad (A \text{ "almost } \cong" B)$$

($X + A \cong X + B$ symmetric)

$$nX \cong nY \quad \text{iff} \quad X \cong Y \quad (n \text{ left cancels})$$

$$nX \cong mX \quad (\text{dichotomy})$$

$$nX \cong mY \quad (\text{can cancel and divide})$$

$$X + Y \cong Y + X \quad (\text{can characterize such pairs})$$

Arithmetic of (LO, \times)

...what about the corresponding isomorphisms for (LO, \times) ?

Arithmetic of (LO, \times) : questions

$AX \cong X$ iff $X \cong A^\omega \times R$? (A “almost \cong ” 1 ?)

$XA \cong X$ symmetric ?

$AX \cong BX$ iff (A “almost \cong ” B ?)

$XA \cong XB$ symmetric ?

$X^n \cong Y^n$ iff $X \cong Y$? (can take n -th roots ?)


$X^n \cong X^m$ (dichotomy ?)

$X^n \cong Y^m$ (Euclidean in exponent ?)

$XY \cong YX$ (can we characterize ?)

Left absorption

Consider the isomorphism $AX \cong X$.



Are there examples where $A \not\cong 1$?

Yes! For an arbitrary A , $X = A^\omega$ works.

Left absorption

Many familiar orders have the form A^ω :

- i. $2^\omega \cong$ the Cantor set,
- ii. $\mathbb{Z}^\omega \cong$ the irrationals,
- iii. $\omega^\omega \cong$ the non-negative reals,
- iv. $\mathbb{Q}^\omega \cong$ the usual example of a G_δ -set.

Left absorption

More generally, if R is arbitrary and if $X \cong A^\omega \times R$ then $AX \cong X$.


Is this general? Not quite!

Thm (E.) $AX \cong X$ iff X is of the form $A^\omega(I_{[u]})$.

... where $A^\omega(I_{[u]})$ denotes a “replacement of A^ω up to tail equivalence” (whatever that means).

Right cancellation

Now consider $AX \cong BX$.



We can't always cancel X , but just like in the additive case, absorption is only barrier!

Thm (Morel): If $AX \cong BX$ then either $A \cong B$ or there is an order $K \not\cong 1$ s.t. $KX \cong X$.

Right cancellation

If $AX \cong BX$, is there a sense in which A is always “almost isomorphic” to B ?

Thm (E. + Paul): Suppose X is a linear order.

- i. For any linear order A , the rule $a \sim_X a' \Leftrightarrow [a, a'] \times X \cong X$ defines a convex equivalence relation on A .
- ii. If $AX \cong BX$, then $A / \sim_X \cong B / \sim_X$.

Right absorption

Now consider the isomorphism $XA \cong X$.

This is *not* symmetric with $AX \cong X$:

- ▶ $AX \cong X$ says “ X can be split into A -many copies of itself.”
- ▶ $XA \cong X$ says “ X can be split into itself-many copies of A .”


We've seen examples: e.g. $\mathbb{Z}2 \cong \mathbb{Z}$.

Can we characterize all examples? Yes (E., unpublished), but more difficult to describe than on other side.

One issue: A^{ω^*} can't be lex-ordered.

Left cancellation

Consider the isomorphism $XA \cong XB$:



Is the left-sided version of Morel's theorem true?

Question: Suppose $XA \cong XB$. Is it true that either $A \cong B$ or there is $L \not\cong 1$ s.t. $XL \cong X$?

Answer (E. + Paul): No. There are even countable counterexamples.

Left cancellation

Is it hopeless to get a nice cancellation result for $XA \cong XB$?

Not quite! We observed that in our counterexamples, the right-hand factors A and B were always *left*-absorbing.

If we assume A, B are not left-absorbing, we get the theorem we want:

Thm (E. + Paul): Suppose $XA \cong XB$ and neither A nor B is left-absorbing. Then either $A \cong B$ or there is $L \not\cong 1$ s.t. $XL \cong X$.

We also showed: this is the best possible left-sided version of Morel's theorem.

Non-absorbing right factors

Even better: under the assumption that A, B are not left-absorbing, we can completely analyze four term isomorphism $XA \cong YB$.

Thm (E. + Paul): Suppose $XA \cong YB$ and neither A nor B is left-absorbing. Then:

- i. If neither A nor B convexly embeds in the other, we have $X \cong Y$.
- ii. If B convexly embeds in A but A does not convexly embed in B , then exactly one holds:
 - a. There is an infinite linear order L s.t. $A \cong LB$ and $Y \cong XL$.
 - b. There are $m, n \in \mathbb{N}$, $m \neq n$, and a linear order C such that $A \cong mC$, $B \cong nC$, and $Xm \cong Yn$.
- iii. If A and B are convexly bi-embeddable, then $X \cong Y$.

Absorbing right factors

So what if the right-hand factors A, B in the isomorphism $XA \cong YB$ are left-absorbing?

We conjecture: if we mod out the lefthand factors by the absorption relations \sim_A, \sim_B , we get the theorems we want.

Conj (E. + Paul): Suppose $XA \cong YB$. Then:

- i. If neither A nor B convexly embeds in the other, we have $X \cong Y$.
- ii. If B convexly embeds in A but A does not convexly embed in B , then there is a linear order L s.t. $XL / \sim_B \cong Y / \sim_B$.
- iii. If A and B are convexly bi-embeddable, then the condensations \sim_A and \sim_B coincide, and we have then $X / \sim \cong Y / \sim$.

Absorbing right factors


Here is the corresponding conjecture for the isomorphism $XA \cong XB$.

Conj (E. + Paul): Suppose $XA \cong XB$. Then either $A \cong B$, or there is an order L such that $XL/\sim \cong X/\sim$.

Says: X is left-cancelling iff X is non-right-absorbing (up to the condensation induced by left-absorption of the right-hand factors).

Taking square roots

Consider the isomorphism $X^2 \cong Y^2$:



Does it follow $X \cong Y$?

Thm (Morel, Sierpinski): No. There exist countable orders $X \not\cong Y$ s.t. $X^2 \cong Y^2$.

Taking square roots

However, in all known cases of $X^2 \cong Y^2$, X and Y are convexly bi-embeddable (i.e. “extremely close” to being isomorphic).

Question: Is this always the case?

Thm (E. + Paul) Yes for countable X, Y .

Morel and Sierpinski's example

The orders X, Y that Morel and Sierpinski constructed have the property that $X \not\cong Y$ but $X^n \cong Y^n \cong Y$ for all $n \geq 2$.

Question (Sierpinski): Does $X^n \cong Y^n$ for some $n > 2$ imply $X^2 \cong Y^2$?

Conj (E. + Paul) Yes for countable X and Y .

Conj (E.) No in general.

Power dichotomy

For a linear order X , is it true that the finite powers X^n , $n \geq 1$ are either all isomorphic or all distinct?

Thm (Morel and Sierpinski): No.


Their example gives X s.t. $X^2 \cong X^3 \cong \dots$ but $X \not\cong X^2$.

However, we do have the following weaker dichotomy:

Thm (E.) $X \cong X^n$ for some $n > 1$ iff $X \cong X^n$ for all $n > 1$.

Commuting pairs

Consider the isomorphism $XY \cong YX$:



Two “obvious” ways it can hold:

- i. (finite product) $\exists C$ s.t. $X \cong C^n$ and $Y \cong C^m$ for some $m, n \in \mathbb{N}$,
- ii. (bi-absorption) $XY \cong YX \cong Y$.

Commuting pairs

Question: Are there multiplicative analogues X, Y of Lindenbaum's “irrational rotation” additive commuting pairs?

Question: If so, are these the only three possible types of multiplicatively commuting pairs X, Y ?

Thank you!